

NOTES AND COMMENTS



Effect of genotype and environment on parasite and pathogen levels in one apiary - a case study

Roy M Francis¹, Esmail Amiri¹, Marina D Meixner^{2*}, Per Kryger¹, Anna Gajda³, Sreten Andonov⁴, Aleksandar Uzunov⁴, Grazyna Topolska³, Leonidas Charistos⁵, Cecilia Costa⁶, Stefan Berg⁷, Malgorzata Bienkowska⁸, Maria Bouga⁹, Ralph B uchler², Winfried Dyrba¹⁰, Fani Hatjina⁵, Evgeniya Ivanova¹¹, Nikola Kezic¹², Seppo Korpela¹³, Yves Le Conte¹⁴, Beata Panasiuk⁷, Hermann Pechhacker¹⁵, George Tsoktouridis¹⁶ and Jerzy Wilde¹⁷

¹Department of Agroecology, University of  rhus, Flakkebjerg, 4200 Slagelse, Denmark.

²LLH, Bee Institute, Erlenstrasse 9, 35274 Kirchhain, Germany.

³Warsaw University of Life Sciences, Faculty of Veterinary Medicine, Department of Pathology and Veterinary Diagnostics, Laboratory of Bee Diseases, Ciszewskiego 8 Street, 02-787 Warsaw, Poland.

⁴Faculty of Agricultural Sciences and Food, bul. Aleksandar Makedonski b. b., 1000 Skopje, Republic of Macedonia.

⁵Hellenic Institute of Apiculture – Hellenic Agr. Org. 'DEMETER', N. Moudania 63200, Greece.

⁶Consiglio per la Ricerca e la sperimentazione in Agricoltura – Unit  di ricerca di apicoltura e bachicoltura (CRA-API), Via di Saliceto 80, 40128 Bologna, Italy.

⁷Bayerische Landesanstalt f r Weinbau und Gartenbau, Bee Division, An der Steige 15, 97209 Veitsh chheim, Germany.

⁸Research Institute of Horticulture, Apiculture Division, 24-100 Puławy, Poland.

⁹Agricultural University of Athens, Laboratory of Agricultural Zoology and Entomology, 75 Iera Odos St., Athens 11855 Greece.

¹⁰Landesverband der Imker Mecklenburg und Vorpommern e.V., Bee Breeding Centre Bantin, Wittenburger Stra e 3, 19246 Bantin, Germany.

¹¹University of Plovdiv, "Paisii Hilendarski", Department of Developmental Biology, Section of Genetics, 24 Tsar Asens St, Plovdiv 4000, Bulgaria.

¹²Faculty of Agriculture, University of Zagreb, Svetosimunska 25, 10000 Zagreb, Croatia.

¹³MTT, Agrifood research Finland, 31600 Jokioinen, Finland.

¹⁴INRA, UR 406 Abeilles et Environnement, Laboratoire Biologie et Protection de l'abeille, Site Agroparc, 84914 Avignon, France.

¹⁵Austrian Carnica Association, Sulzbach 1, 3293 Lunz am See, Austria.

¹⁶Laboratory of Conservation and Evaluation of Native and Floricultural Species – Hellenic Agr. Org. 'DEMETER', PO Box 60125, 57 001 Thermi-Thessaloniki, Greece.

¹⁷Apiculture Division, Warmia and Mazury University, Sloneczna 48, 10-710 Olsztyn, Poland.

Received 13 January 2014, accepted subject to revision 13 March 2014, accepted for publication 28 April 2014.

*Corresponding author: Email: marina.meixner@llh.hessen.de

Keywords: COLOSS, Genotype-Environment Interactions Experiment, *Apis mellifera* L., honey bee, *Varroa destructor*, *Nosema* spp., quantitative virus analysis, survival

An international collaborative experiment was run from 2009 to 2012 (Costa *et al.*, 2012) with the aim of understanding genotype-environment effects on survival and health status of honey bee colonies headed by queens of different European origins that were tested in various locations under differing environmental conditions. No chemical treatment against bee diseases was performed on any of the test colonies. The colonies were managed according to a common protocol, which included an assessment of their health status (for details, see Meixner *et al.*, 2014). One apiary located in Chalkidiki, Greece, was selected for a detailed case study analysis of parasites and pathogens, including an absolute quantification of viral titres.

The colonies were sampled regularly between autumn 2009 and autumn 2011 and analysed for *Varroa destructor* and *Nosema* spp. as described in Meixner *et al.* (2014). For virus analysis, approximately

30 worker bees from each colony were sampled in March and September of 2010, and April and October of 2011, and analysed for deformed wing virus (DWV), black queen cell virus (BQCV), acute bee paralysis virus (ABPV), Kashmir bee virus (KBV) and Israeli paralysis virus (IAPV). Extraction of RNA, qPCR and quantification of virus titres followed the methods described in Francis *et al.* (2013), with ABPV, KBV and IAPV assessed together in a single assay (AKI). Viral titres (acute paralysis complex (AKI), DWV and BQCV), *Nosema* spp. infection levels and *V. destructor* mite infestation levels during the seven sampling times were compared to the survival status of the colony at the end of the experiment. The data were analysed for patterns, trends, correlations and significant differences.

Of the 39 starting colonies, 31 could be sampled for quantitative virus analysis in the spring of 2010. They belonged to the genotypes




Fig. 1. Overview of results at one location (Chalkidiki, Greece). Viral titres (ABPV-KBV-IAPV (AKI), DWV and BQCV), *Nosema* spp. spore levels and *V. destructor* mite levels are shown for four genotypes (CarV, LigI, MacB and MacG (local bee)) at seven sampling periods. Survival status was determined in April every year, and colonies that remained alive after the experiment are marked 'Alive'. Viruses were sampled in spring and autumn of 2010 and 2011. *Nosema* spp. were sampled in three seasons of the years 2010 and 2011. *V. destructor* was sampled at the same times as *Nosema* spp., and also in the autumn of 2009. Note that not all colonies were sampled for all pathogens for all time periods. MacG colonies showed the lowest mite levels and DWV titres in the autumn of 2010 and 2011. The local bee (MacG) had the highest number of surviving colonies (6 of 9) at the end of the experiment.

CarV (*A. m. carnica*; n = 6), LigI (*A. m. ligustica*; n = 6), MacB (*A. m. macedonica*; n = 10) and MacG (*A. m. macedonica*; n = 9). The genotype of local origin was MacG (for details of genotypes see Francis *et al.*, 2014). At the end of the experiment, eight colonies survived, including MacG (n = 6), MacB (n = 1) and LigI (n = 1). None of the *A. m. carnica* colonies survived the second winter (2010/11).

V. destructor infestation levels generally increased over the duration of the experiment, but the local bees (MacG) showed significantly lower mite infestation rates in autumn of 2009 and 2010 compared to the three other genotypes (GLM, p < 0.05) (see Fig. 1.). Mite levels in October 2010 and August 2011 were negatively correlated to the survival duration of the colonies in days ($R^2 = 0.63$, p < 0.001).

The viral titres of the colonies and showed a strong seasonal trend (see Fig. 1.). In both years, the ABPV complex and DWV showed low titres in spring, and high titres in autumn, with a significant difference (Wilcoxon, $P < 0.05$) between them. DWV showed overall higher titres than AKI. While the titres of AKI in October 2010 were highly variable, the infection with DWV was homogeneously high. In 2010 and 2011, the number of *V. destructor* mites in autumn significantly correlated with DWV titres in the spring of the following year ($R^2 = 0.64$, $p < 0.01$). The MacG colonies consistently had the lowest mite infestation and the lowest DWV titres, while DWV titres in non-local bees were significantly higher (Wilcoxon, $W = 37$, $p < 0.05$).

Following Francis *et al.* (2013), we consider bees harbouring more than 10^7 copies of virus to represent an overt virus infection. In the case of DWV infection, we observe an increase in titres greater than 10^7 copies from 7% of the colonies in April ($n = 15$), to 55% of the colonies in September ($n = 11$) and 78% of the colonies in October ($n = 23$). In contrast, only one colony out of 23 had AKI titres greater than 10^7 copies in October.

Nosema spp. infection levels were higher in spring and autumn than in the summer (Fig. 1.). In spring 2010, all colonies had less than 5 million *Nosema* spp. spores per bee, except two *A. m. ligustica* colonies that harboured more than 15 million spores (significantly higher, GLM, $p < 0.05$). One of these colonies eventually died, while the other survived. A similar trend in *Nosema* spp. infection was observed in spring 2011. One LigI colony survived in spite of an extremely high *Nosema* spp. level. The BQCV infection level showed a low-high-low trend, with low titres at the beginning (spring 2010), high titres in the middle (autumn 2010 - spring 2011) and low titres towards the end (autumn 2011) of the experiment. The highest BQCV titres were observed parallel to elevated *Nosema* spp. levels in the same colonies in spring and autumn. In spring 2011, *Nosema* spp. levels were significantly correlated to BQCV titres ($R^2 = 0.68$, $p < 0.01$). In spring 2010, for example, LigI colonies had the highest *Nosema* spp. spore loads as well as the highest BQCV titres.

Although our data do not allow us to deduce a causal relationship between mortality and virus infections in this apiary, the patterns of infection differ conspicuously among introduced and local bees. A similar pattern can be observed for the infection with *Nosema* spp., with the *A. m. ligustica* origin showing the highest levels, also reflected in comparatively high BQCV titres in the same colonies. No clear patterns of survival or death of colonies in relation to *Nosema* spp. or BQCV titres emerge from these results.

Given the close proximity of the colonies in a single apiary, our findings are surprising, since a particular protection of the colonies of local origin from pathogen transfer between hives appears unlikely. Rather, our results suggest that bees of local origin are better in tune with environmental factors related to flowering patterns, climatic variation and locally prevailing apicultural methods and therefore may command more sufficient resources to fend off pathogens.

Acknowledgements

The Genotype and Environment Interactions experiment and this Special Issue of the *Journal of Apicultural Research* were conceived at meetings and workshops organised by COLOSS. The COLOSS (Prevention of honey bee COLony LOSSes) association (<http://coloss.org/>) is an international, non-profit association based in Bern, Switzerland that is focussed on improving the well-being of bees at a global level. Between 2008 and 2012, COLOSS was funded by COST (European Cooperation in Science and Technology) through the COST Action FA0803. We gratefully acknowledge COLOSS, not only for funding numerous workshops during the course of the experiment which facilitated the exchange of samples and ideas, and the analyses described in this paper, but also for the excellent collaboration and warm working atmosphere. COLOSS is now supported by the Ricola Foundation - Nature & Culture and has funded Open Access for this paper.

References

- COSTA, C; BERG, S; BIENKOWSKA, M; BOUGA, M; BUBALO, D; BÜCHLER, R; CHARISTOS, L; LE CONTE, Y; DRAZIC, M; DYRBA, W; FILLIPI, J; HATJINA, F; IVANOVA, E; KEZIC, N; KIPRIJANOVSKA, H; KOKINIS, M; KORPELA, S; KRYGER, P; LODESANI, M; MEIXNER, M; PANASIUK, B; PECHHACKER, H; PETROV, P; OLIVERI, E; RUOTTINEN, L; UZUNOV, A; VACCARI, G; WILDE, J (2012) A Europe-wide experiment for assessing the impact of genotype-environment interactions on the vitality of honey bee colonies: methodology. *Journal of Apicultural Science* 56: 147-158. <http://dx.doi.org/10.2478/v10289-012-0015-9>
- FRANCIS, R M; KRYGER, P; MEIXNER, M; BOUGA, M; IVANOVA, E; ANDONOV, S; BERG, S; BIENKOWSKA, M; BÜCHLER, R; CHARISTOS, L; COSTA, C; DYRBA, W; HATJINA, F; PANASIUK, B; PECHHACKER, H; KEZIĆ, N; KORPELA, S; LE CONTE, Y; UZUNOV, A; WILDE, J (2014b) The genetic origin of honey bee colonies used in the Genotype-Environment-Interactions experiment: a comparison of methods. *Journal of Apicultural Research* 53(2): 188-204. <http://dx.doi.org/10.3896/IBRA.1.53.2.02>
- FRANCIS, R M; NIELSEN, S L; KRYGER, P (2013) Patterns of viral infection in honey bee queens. *Journal of General Virology* 94(3): 668-676. <http://dx.doi.org/10.1099/vir.0.047019-0>
- MEIXNER, M D; FRANCIS, R M; GAJDA, A; KRYGER, P; ANDONOV, S; UZUNOV, A; TOPOLSKA, G; COSTA, C; AMIRI, E; BERG, S; BIENKOWSKA, M; BOUGA, M; BÜCHLER, R; DYRBA, W; GURGULOVA, K; HATJINA, F; IVANOVA, E; JANES, M; KEZIC, N; KORPELA, S; LE CONTE, Y; PANASIUK, B; PECHHACKER, H; TSOKTOURIDIS, G; VACCARI, G; WILDE, J (2014) Occurrence of parasites and pathogens in honey bee colonies used in a European genotype - environment - interactions experiment. *Journal of Apicultural Research* 53(2): 215-229. <http://dx.doi.org/10.3896/IBRA.1.53.2.04>